

Review

Using 2,4-dinitrophenylhydrazine in spectrophotometric determination

Safaa Abdulaleem Zakaria*, Zena Talal and Nabeel Sabeeh Othman

Chemistry Department, College of Science, University of Mosul, Mosul, Iraq

<https://doi.org/10.54153/sjpas.2022.v4i2.368>

Article Information

Received: 04/03/2022

Accepted: 09/05/2022

Keywords:

2,4-Dinitrophenylhydrazine,
Aldehyde and ketones,
Spectrophotometric method,
Oxidative coupling,
diazotization-coupling

Corresponding Author

E-mail:

nsn20002004@uomosul.edu.iq

Mobile: 07703379110

Abstract

2,4-Dinitrophenylhydrazine (2,4-Dnphh) is substituted hydrazine. It is structure contains hydrazine which is the main active functional group. 2,4-Dnphh is an important reagent in analytical chemistry it is used firstly in the identification (qualitative analysis) of carbonyl groups (aldehyde and ketones compounds). 2,4-Dnphh is react with aliphatic and aromatic carbonyl compounds to produce the corresponding 2,4-dinitrophenyl-hydrazone. The aromatic carbonyls give red precipitates compared with aliphatic carbonyls giving more yellow color. The two colored products were achieved according to the addition-elimination reaction, the reaction included nucleophilic addition of amino group (-NH₂) to the carbonyl group(C=O), then the elimination of the H₂O molecule. Also, a modification of 2,4-Dnphh to produce a derivative can be used as a chemical sensor for some colorimetric detection of various compounds. Recently 2,4-Dnphh is used in quantitative spectrophotometric estimation of various compounds via oxidative coupling reaction and diazotization-coupling reaction, and also 2,4-dinitrophenylhydrazone-compound prepared from the reaction of 2,4-Dnphh with compound to be determined give good results in injection to the column of high-performance liquid chromatography. The present review included all methods as possible in the literature including the uses of 2,4-Dnphh in the qualitative and quantitative determination of various compounds.

Introduction:

2,4-Dinitrophenylhydrazine is one of the important organic compounds has the chemical formula [C₆H₃(NO₂)₂NHNH₂]. 2,4-Dnphh is a solid compound with orange-red color, it is chemical structure (Figure 1) contains two nitro groups in ortho and para positions to hydrazine so it is substituted hydrazine.





Fig. 1: The chemical structure of 2,4-Dnphh.

2,4-Dnphh can be prepared by the reaction of 2,4-dinitrochlorobenzene with hydrazine sulphate in the presence of $\text{N}_2\text{H}_4\cdot\text{H}_2\text{SO}_4$ and KOAc (Scheme 1)[1].




Scheme 1: Preparation of 2,4-Dnphh.

Dissolving 2,4-Dnphh in aqueous solution to prepared Brady's reagent, to be used for the identification of carbonyl compounds (aldehyde and ketone).

Other researchers prepared Brady's reagent or Borche's reagent, by dissolving 2,4-dinitrophenylhydrazine in a mixture solution containing methanol and a few milliliters of concentrated sulfuric acid[2].


The formation of a yellow, orange or red precipitate is indicated as a positive test of the presences of carbonyl groups. The aromatic carbonyls give red precipitates compared with aliphatic carbonyls give a more intense yellow color.[3]

2,4-Dnphh can reacts with aldehydes and also with ketones to form a 2,4-dinitrophenylhydrazone. An example reaction with ethanal [4] (Scheme 2).



Scheme 2: Reaction of 2,4-Dnphh with ethanal.


The condensation reaction to form the product (hydrazone) from the reaction of ketone with 2,4-Dnphh included loss of water, this suggested mechanism can be considered as an example of addition-elimination reaction, of the amino group (-NH2) undergo nucleophilic addition to the carbonyl group(C=O), then the elimination of a water molecule, as per the following equation:[5].



Recently a modification of the derivative 2,4- dinitrophenylhydrazine dithiocarbamate - silver nanoparticles can be used as am chemical sensor for some colorimetric detection of various compounds such as omeprazole [6].


The liquid carbonyl compounds, for example, cyclohexanone can react with 2,4-Dnphh to produce solid derivatives. The reaction happens at one of the two NH_2 groups. The ion-pair

electrons of the NH_2 group attached to the carbonyl is less nucleophilic compared to those of the other NH_2 group (Scheme 3) [7].




Scheme 3: The suggested reaction of cyclohexanone with 2,4-Dnphh.

2,4-Dnphh has been often used as a derivatizing agent for carbonyl compounds, including reducing compounds for example glucose to form 2,4-Dnphh- glucose derivative (Scheme 4), acetaldehyde, and various carbonyl groups, and these derivatives can easily determination by HPLC techniques. [8-10].




Scheme 4: 2,4-Dnphh- glucose derivative.

It has been shown that 2,4-Dnphh can be oxidized by various oxidizing agents such as potassium iodate or sodium periodate to produce diazotized 2,4- Dnphh as an intermediate compound which later can be under go coupling reaction with other compounds such as pyridoxine hydrochloride in alkaline medium to produce red azo dye that has the highest absorption at the wavelength of 522 nm. The proposed reactions as shown below in Scheme 5 [11].



Scheme 5: The proposed reactions to produce the red azo dye
The suggested structure of red azo dye is shown below.



Through the literature survey, many researchers used 2,4- Dnphh reagent in the estimation of various compounds. The reaction mechanism was explained concerning oxidation and coupling reaction, relying mainly on the use of the oxidizing agent in the first step of the reaction, and below Table 1 includes these types of reactions

Table 1: Using 2,4- Dnphh in determination various compound via oxidative coupling reaction.

Compound	λ_{max} ,nm	Linearity μg/ml	ϵ L/mol.cm	Reference
Fungicide dithianon	550	[12]
Selenium(IV)	520	0.03-3.5	3.1×10^4	[13]
Vanadium	495	0.02-3.5		[14]
Vitamin C	280	0.0323	[15]
Atorvastatin	479	4-10	558.65×10^3	[16]
Ezetimibe	457	4-12	409.34×10^3	
Acyclovir&	414	20-60	[17]
Ganciclovir	450	5-25		
Modafinil	476	2-7	3.3297×10^5	[18]
Carbamazepine	485	4-50	6.7335×10^3	[19]
Bisacodyl	522	2-20	6505	[20]
Mefenamic acid	528	2-8	3.6873×10^4	[21]
Sulphadiazine	486	3-15	2.23×10^4	[22]
Propranolol hydrochloride	481	1-35	6224.16	[23]
Catechol	385	1-40	5.0866×10^3	[24]
Resorcinol	404	1-25	4.404×10^3	
Lansoprazole	484.5	1-30	13260.132	[25]
Nateglinide	480	30-100	0.331054×10^3	[26]
Pregabalin	461	50-450	0.273824×10^3	[27]
Clopidogrel	412	10-30	0.8817×10^4	[28]
Gemfibrozil	421	2-10	2.6081×10^4	

While Table 2 also includes the estimation of various compounds, the researchers suggested an explanation for the reaction based on the principle of diazotization and coupling, depending on the interpretation of the formation of corresponding diazonium salt of 2,4-Dnphh when the reagent was oxidized by the oxidizing reagents.

Table 2: Using 2,4- Dnphh in determination various compounds via diazotization and coupling reaction.

Compound	$\lambda_{\text{max}}, \text{nm}$	Linearity $\mu\text{g/ml}$	ϵ L/mol.cm	Reference
Salbutamol	540	2.5-17	2.19×10^4	[29]
Ritodrine	510	2-29	0.74×10^4	
Amoxicillin Isoxsuprine	520	4-33	1.00×10^4	
	520	5-30	8.04×10^4	
Thymol	570	0.25-10	2.2×10^4	[30]
Diclofenac sodium	600	0.5-40	2.0994×10^4	[31]
Cefadroxil	515	7.5-75	7×10^3	[32]
Oxymetazoline hydrochloride	607	2.5-30	7242	[33]
Sulfathiazole	492	2-28	1.1437×10^4	[34]
Paracetamol	430	0.5-18.4	2.776×10^4	[35]
		mg.mL ⁻¹		
Tetracycline hydrochloride	360	0.1-9	1.262×10^4	[36]
Dopamine	560	5-50	7.9×10^4	[37]
Sulphamethoxazole	435	2-14	14412.77	[38]

Table 3 included the compounds determined via condensation reaction with 2,4-Dnphh to produce the corresponding hydrazone.

Table 3: Using 2,4- Dnphh in determination various compounds via condensation reaction.

Compound	$\lambda_{\text{max}}, \text{nm}$	Linearity $\mu\text{g/ml}$	ϵ L/mol.cm	Reference
Josamycin	411	10-160	[39]
Eplerenone	430	5-35	3.177×10^4	[40]
Sitagliptin phosphate	400	2-10	0.2485×10^{-4}	[41]
Cilostazol	355	2-20	2.2980×10^4	[42]
Haloperidol	456	3.6-10.8	2.84×10^4	[43]
Artemisinin	310	0.5-60	2.930×10^4	[44]
Dihydroartemisinin	300	0.5-50	2.8×10^4	
Macrolide antibiotics	542-545	5-40	1.48×10^4	[45]
	523-526	5-35	2.59×10^4	
	539-542	5-35	2.10×10^4	
Tolperrisone	520	2.5-15	2.4×10^4	[46]
Lamivudine	438	5-35	[47]
Reactive carbonyl in proteins and peptides	370	[48]

Conclusion

2,4-Dinitrophenylhydrazine is an important reagent used to identify(qualitative) of carbonyl groups (aldehyde and ketones compounds), and also it is used in quantitative spectrophotometric determination of various compounds via various types of reactions, such as oxidative coupling, diazotization and coupling reaction and reacted with aliphatic and aromatic carbonyl compounds to produce the corresponding 2,4- dinitrophenyl-hydrazone (addition-elimination) reactions.

References

- 1-Allen, C. (1933). "2,4-Dinitrophenylhydrazine". *Organic Syntheses*. 13, 36.
2. Jddles, H. A., & Jackson, C. E. (1934). Determination of carbonyl compounds by means of 2, 4-dinitrophenylhydrazine. *Industrial & Engineering Chemistry Analytical Edition*, 6(6), 454-456. <https://doi.org/10.1021/ac50092a030>
- 3.<http://wiki.colby.edu/download/attachments/110920618/Experiment+%232.pdf?version=1&modificationDate=1265312071267>
4. <https://byjus.com/chemistry/2-4-dinitrophenylhydrazine/#Reaction-with-Ethanal>
5. Lawrence, R. C. (1965). Use of 2, 4-dinitrophenylhydrazine for the estimation of micro amounts of carbonyls. *Nature*, 205(4978), 1313-1314.
6. Jabeen, S., Shah, M. R., Rauf, A., Rauf, S., Jabbar, A., & Bhanger, M. I. (2017). 2, 4-Dinitrophenylhydrazine dithiocarbamate modified silver nanoparticles as sensor for the colorimetric determination of omeprazole. *Current Analytical Chemistry*, 13(6), 532-539. DOI: <https://doi.org/10.2174/1573411013666170911155757>
7. Ouellette, R. J., & Rawn, J. D. (2014). 1-structure and bonding in organic compounds. *Organic Chemistry*, 1-39.
8. Guan, X., Rubin, E., & Anni, H. (2012). An optimized method for the measurement of acetaldehyde by high-performance liquid chromatography. *Alcoholism: Clinical and Experimental Research*, 36(3), 398-405. <https://doi.org/10.1111/j.1530-0277.2011.01612.x>
9. Barman, B. N. (2014). Accurate determination of aldehydes in amine catalysts or amines by 2, 4-dinitrophenylhydrazine derivatization. *Journal of Chromatography A*, 1327, 19-26. <https://doi.org/10.1016/j.chroma.2013.12.064>
10. Uchiyama, S., Inaba, Y., & Kunugita, N. (2011). Derivatization of carbonyl compounds with 2, 4-dinitrophenylhydrazine and their subsequent determination by high-performance liquid chromatography. *Journal of Chromatography B*, 879(17-18), 1282-1289. <https://doi.org/10.1016/j.jchromb.2010.09.028>
- 11- Saleem, M. S., & Mohammed, Z. A. (2013). A spectrophotometric micro determination of pyridoxine hydrochloride by coupling diazometry. *JOURNAL OF EDUCATION AND SCIENCE*, 26(5), 23-35. DOI:[10.33899/edusj.2013.163052](https://doi.org/10.33899/edusj.2013.163052)
12. Yuen, S. H. (1969). Spectrophotometric determination of the fungicide dithianon in aqueous formulation Delan-Col with 2, 4-dinitrophenylhydrazine. *Analyst*, 94(1125), 1095-1098. <https://doi.org/10.1039/AN9699401095>
13. Krishnaiah, L., Kumar, K. S., Suvardhan, K., & Chiranjeevi, P. (2003, December). Simple spectrophotometric determination of traces of selenium in environmental samples. In *Proceedings of the third international conference on environment and health, Chennai, India* (pp. 15-17).

14. Priya, B. K., Subrahmanyam, P., Kumar, J. D., & Chiranjeevi, P. (2006). Simple sensitive spectrophotometric determination of vanadium in biological and environmental samples. *E-Journal of Chemistry*, 3(4), 286-297.
15. Hamad, Q. Y. M. M. (2009). Spectrophotometric determination of total vitamin c in some fruits and vegetables at Koya area-Kurdistan region/ Iraq. *Kirkuk university journal for scientific studies*, 4(2),46-54.
16. Shravya, A., Ch, R. S., Gurupadayya, B. M., & Sireesha, M. (2010). Spectrophotometric determination of atorvastatin and ezetimibe using 2, 4-DNP in bulk and pharmaceutical dosage forms. *IJPPT* , 2 (4)1046-1056.
17. Anil Kumar, T., Gurupadayya, B. M., Reddy, R., & PrudhviRaju, M. V. (2012). Selective and validated Spectrophotometric methods for determination of acyclovir and ganciclovir with 2, 4 as reagent. *Journal of Applied Chemical Research*, 6(1), 14-24.
18. Rashmi, N. G., Chandan, R. S., Gurupadayya, B. M., Srujana, S., & Raagaleena, V. (2012). Validated analytical determination of modafinil by 2, 4-DNP and PDAC reagents. *Journal of Pharmacy Research*, 5(5), 2527-2531.
19. Fadhel S. R., Abdulla N. I. Sulaiman I. D. (2016). Spectrophotometric determination of carbamazepine via oxidative coupling reaction with 2,4-dinitrophenyl hydrazine. *Ibn Al-Haitham Jour. for Pure & Appl. Sci.*29(1).226-238.
20. Mohammed, I. S., Nasser, K. A., & Mhemeed, A. H. (2017). Spectrophotometric determination of bisacodyl in pure and pharmaceutical preparation via oxidative coupling organic reaction. *Baghdad Science Journal*, 14(1),181-188. DOI: <http://dx.doi.org/10.21123/bsj.2017.14.1.0181>
21. Bakir, M. H. (2017). Spectrophotometric Determination mefenamic acid, in pharmaceutical preparation by oxidative coupling 2, 4, dinitrophenyl hydrazine by N-bromo succinimide. *Kirkuk University Journal-Scientific Studies*, 12(3), 649-666. DOI: 10.32894/kujss.2017.131503
22. Ahmed, N. A. A. S., & Khaleel, A. I. (2018). Spectrophotometric determination of sulphadiazine Using 2, 4-dinitrophenylhydrazine as Coupling Reagent. *Tikrit Journal of Pure Science*, 23(8), 77-82. <<http://tjps.tu.edu.iq/index.php/j/article/view/152>>
23. Habeeb, E. D., & Sulaiman, I. D. Spectrophotometric determination of propranolol hydrochloride via oxidative coupling reaction with 2, 4-dinitrophenyl hydrazine. *International Journal of Drug Delivery Technology*,11(1),29-35.
24. Mohammed, D. H., & Omar, F. K. (2021). Spectrophotometric determination of catechol and resorcinol by oxidative coupling With 2, 4-dinitrophenyl Hydrazine. *Egyptian Journal of Chemistry*, 64(9), 5061-5065. DOI: 10.21608/EJCHEM.2021.57506.3253
25. Jamel, S. H., & Sulaiman, I. D. Spectrophotometric estimation of the Lansoprazole by oxidative coupling reaction with 2, 4-dinitrophenyl hydrazine. *IJDDT*, 11 (I) ,52-57. DOI: 10.25258/ijddt.11.1.9
26. Sireesha, M., Chandan, R. S., Gurupadayya, B. M., & Shravya, A. (2011). Spectrophotometric determination of Nateglinide using 2, 4-dinitrophenyl hydrazine and potassium ferricyanide in pharmaceutical dosage form. *Der Pharma Chemica*, 3(1), 497-06. <http://derpharmacemica.com/archive.html>.
27. Sowjanya, K., Thejaswini, J. C., Gurupadayya, B. M., & Indu Priya, M. (2011). Spectrophotometric determination of pregabalin using 1, 2-naphthaquinone-4-sulfonic

acid Sodium and 2, 4 dinitrophenyl hydrazine in pharmaceutical dosage form. *Der Pharmacia Lettre*, 3(2), 47-56. <http://scholarsresearchlibrary.com/archive.html>

28. Srujana, S., Chandan, R. S., Gurupadayya, B. M., Raagaleena, V., & Manoja, K. M. L. (2012). Validated spectroscopic determination of clopidogrel and gemfibrozil using 2, 4-dinitro phenyl hydrazine reagent. *J Pharm Res*, 5(4), 1906-1909.

29 . Nagaraja, P., & Shrestha, A. K. (2010). Spectrophotometric method for the determination of drugs containing phenol group by using 2, 4-dinitrophenylhydrazine. *E-Journal of Chemistry*, 7(2), 395-402. <https://doi.org/10.1155/2010/328061>

30. Fadhil, G. (2014). Spectrophotometric determination of thymol in pharmaceutical preparations via oxidative coupling reaction with 2, 4-dinitrophenylhydrazine in the Presence of Potassium Periodate. *Iraqi J. Sci*, 55(1), 27-34.

31. Dikran, S. B., & Mahmood, R. M. (2017). Spectrophotometric determination of diclofenac sodium using 2, 4-dinitrophenylhydrazine in pure form and pharmaceutical preparations. *Ibn AL-Haitham Journal For Pure and Applied Science*, 28(3), 129-141.

32. Almasri, I., Ramadan, M. & Khayal, G.(2015). Spectrophotometric determination of cefadroxil in bulk and dosage forms using 2,4-dinitrophenylhydrazine. *Journal of Al Azhar University-Gaza (Natural Sciences)*, 17 ,129-146.

33. Humeidy, I. T. (2018). Spectrophotometric method for Determination of oxymetazoline, HCl in a Pharmaceutical formulation using 2, 4-dinitrophenylhydrazine. *Tikrit Journal of Pure Science*, 20(2), 88-96. <http://tjps.tu.edu.iq/index.php/j/article/view/200>

34. Humedy, I. T., & Ali, I. M. (2017). Spectrophotometric determination of sulfathiazole using 2, 4-dinitrophenylhydrazine as coupling reagent. *Ibn AL-Haitham Journal For Pure and Applied Science*, 29(2), 227-239.

35. Pasha, C. (2020). Determination of paracetamol in pharmaceutical samples by spectrophotometric method. *Eclética Química*, 45(3), 37-46. DOI: <https://doi.org/10.26850/1678-4618eqj.v45.3.2020.p37-46>

36. Khaleel, R. M., & Mohammed, D. H. (2020, November). Spectrophotometric determination of tetracycline hydrochloride using 2, 4-dinitrophenyl hydrazine as coupling reagent. In *Journal of Physics: Conference Series* (Vol. 1664, No. 1, p. 012084). IOP Publishing. J. Phys.: Conf. Ser. ,1-11.

37. Ramadan, M. A. A., Almasri, I., & Khayal, G. (2020). Spectrophotometric determination of dopamine in bulk and dosage forms using 2, 4-dinitrophenylhydrazine. *Turkish Journal of Pharmaceutical Sciences*, 17(6), 679-685. DOI: [10.4274/tjps.galenos.2019.25902](https://doi.org/10.4274/tjps.galenos.2019.25902)

38. Rashidy, A., Mohammed, A. A., Al Badrany, K. A., & Al Garagoly, G. M. (2020). Spectrophotometric determination of sulphamethoxazole drug by new pyrazoline derived from 2, 4-dinitro phenyl hydrazine. In *Materials Science Forum* (Vol. 1002, pp. 350-359). Trans Tech Publications Ltd. <https://doi.org/10.4028/www.scientific.net/MSF.1002.350>

39. Alarfaj, N. A., Razeq, S. A., & Alsehaly, H. M. (2011). Determination of josamycin and ciprofloxacin in their pharmaceutical dosage forms by spectrophotometry. *Asian Journal of Chemistry*, 23(8), 3362

40. Ganure, A. L., Dehghan, M. H. G., Patel, D. S., & Sangshetti, J. N. Spectrophotometric determination of Eplerenone in pharmaceutical bulk and tablet dosage form.. *Der Pharma Chemica*, 3(1) ,420-425.

41. Disha, N. S., & Gurupadhyaya, B. M. (2015). Spectrophotometric determination of Sitagliptin phosphate in bulk and pharmaceutical formulations. *International Journal of Innovative Science, Engineering and Technology*, 2(7), 702-709.

42. Kuruba, S. A., & Hanamshetty, P. C. (2015). A simple spectrophotometric quantitative determination of Cilostazol in bulk and pharmaceutical dosage forms using DNPH reagent. *Journal of Applied Pharmaceutical Science*, 5(12), 117-121. [DOI: 10.7324/JAPS.2015.501220](https://doi.org/10.7324/JAPS.2015.501220)

43. Rahman, N., Sameen, S., & Kashif, M. (2016). Spectroscopic study on the interaction of haloperidol and 2, 4-dinitrophenylhydrazine and its application for the quantification in drug formulations. *Analytical Chemistry Letters*, 6(6), 874-885. <https://doi.org/10.1080/22297928.2016.1265898>

44. Emmanuel, A., Anthony, P.C. , Emmanuel, E., AlnoldI, B. & Bartholomew F. C. (2020). A novel spectrophotometric determination of some artemisinin derivatives in tablet using 2,4-DNPHH reagent . *World Journal of Pharmaceutical Research*, 9(7),137-148.

45. Abdelmageed, O. H. (2007). Development and validation of a spectrophotometric method for the determination of macrolide antibiotics by using 2, 4-dinitrophenylhydrazine. *Journal of AOAC International*, 90(2), 364-371. <https://doi.org/10.1093/jaoac/90.2.364>

46. Praveen, P. S., Anupama, B., Jagathi, V., & Rao, G. D. (2010). Spectrophotometric determination of Tolperisone using 2, 4-dinitrophenylhydrazine reagent. *Int J Res Pharm Sci*, 3, 317-320. <http://scopeindex.org/handle/sc/94>

47. Musa, A. (2014). A Novel Spectrophotometric method for determination of lamivudine in pure and tablet dosage form by. *Research & Development*, 6(1), 1-5.

48. Fields, R., & Dixon, H. B. (1971). Micro method for determination of reactive carbonyl groups in proteins and peptides, using 2, 4-dinitrophenylhydrazine. *Biochemical Journal*, 121(4), 587-589. [DOI: 10.1042/bj1210587](https://doi.org/10.1042/bj1210587)

استخدام 2،4-ثنائي نيتروفينيل هيدرازين في التقدير الطيفي

صفاء عبد العليم زكريا، زينه طلال، نبيل صبيح عثمان

قسم الكيمياء، كلية العلوم، جامعة الموصل

الخلاصة:

يعد 2،4-ثنائي نيتروفينيل هيدرازين من معوضات الهيدرازين. إذ يحتوي في تركيبه على الهيدرازين وهو المجموعة الوظيفية الرئيسية النشطة. وبعد كشف مهم في الكيمياء التحليلية اذ تم استخدامه أولاً في التخخيص النوعي لمجموعات الكاربونيل (مركبات الألديهيد والكيتونات). وذلك بتفاعله مع مركبات الكربونيل الأليفاتية والاروماتية لإنتاج 2،4-ثنائي نتروفينيل-هيدرازون المقابل. تعطي مركبات الكاربونيل الاروماتية روابط حمراء مقارنة مع مركبات الكربونيل الأليفاتية التي تعطي على الأغلب اللون الأصفر. تم التحقق من الناتجين ذو اللونين ووجدا انهما يسيران وفقاً لتفاعل الإضافة والحذف، يتضمن التفاعل إضافة المجموعة الأمينية (NH₂) إلى مجموعة الكاربونيل (C = O) ، ثم عملية حذف جزئية ماء. أيضاً يمكن استخدام الكاشف 2،4-ثنائي نيتروفينيل هيدرازين لإنتاج مشتق بوصفه مستشعر كيميائي يستخدم للكشف اللوني للمركبات المختلفة. تم استخدام 2،4-ثنائي نيتروفينيل هيدرازين مؤخراً في التقدير الطيفي الكمي للمركبات المختلفة عن طريق تفاعل الاقتران التاكسي وتفاعل الأزوتة والاقتران، وكذلك 2،4-ثنائي نتروفينيل هيدرازون المحضر من التفاعل مع المركب المراد تحديده يعطي نتائج جيدة في الحقن في عمود كروماتوغرافيا السائل عال الأداء. تضمنت المراجعة الحالية جميع الطرائق المتوفرة في الأدبيات التي تضمنت استخدامات 2،4-ثنائي نيتروفينيل هيدرازين في التقدير النوعي والكمي للمركبات المختلفة.

معلومات البحث:

تأريخ الاستلام: 2022/03/04

تأريخ القبول: 2022/05/09

الكلمات المفتاحية:

2،4-ثنائي نيتروفينيل
هيدرازين، الألديهيدات
وكيتونات، طرائق طيفية، الأكسدة
والاقتران، الأزوتة والاقتران

معلومات المؤلف

الإيميل: nsn20002004@uomosul.edu.iq

الموبايل: 07703379110